

2018

Screening and Identification of Developmental Coordination Disorder in Children with and without ADHD

Shanna Deasley

University of Windsor, deasley@uwindsor.ca

Natalie Gayle Frost

University of Windsor

Lynette Renee Kivisto

University of Windsor

Thomas Anthony Duda

University of Windsor

Follow this and additional works at: <https://scholar.uwindsor.ca/research-result-summaries>

Consistent with the TCPS 2 (4.7) this is a research summary provided to participants and should not be considered a formal publication of results.

Recommended Citation

Deasley, S., Frost, N. G., Kivisto, L. R., & Duda, T. A. (2018). Screening and Identification of Developmental Coordination Disorder in Children with and without ADHD. Retrieved from <https://scholar.uwindsor.ca/research-result-summaries/33>

This Completed Summary is brought to you for free and open access by Scholarship at UWindsor. It has been accepted for inclusion in Research Result Summaries by an authorized administrator of Scholarship at UWindsor. For more information, please contact scholarship@uwindsor.ca.

Title: Screening and Identification of Developmental Coordination Disorder in Children with and without ADHD

Authors: Shanna Deasley, Natalie Frost, Lynette R. Kivisto, Thomas A. Duda, Joseph E. Casey

Objective: Developmental Coordination Disorder (DCD) is characterized by a delay in the acquisition of appropriate motor skills. Motor difficulties associated with DCD relate to problems with handwriting, attention, reading, and social cognition. Individuals with ADHD often have motor skills deficits, and comorbidity with DCD is high (30-50%). Despite this, motor problems and DCD remain under-recognized and undertreated. This study examined whether children with ADHD have more parent reported DCD symptoms than children without ADHD, as well as the prevalence of undiagnosed DCD. It was predicted that children with ADHD would have more DCD symptoms than children without ADHD. As DCD is under-recognized, it was predicted that some children without a DCD diagnosis would screen positive for DCD.

Participants and Methods: Parents ($N=32$) of children ($M_{age}=11.6$ years; 59% male; 50% ADHD) completed the Developmental Coordination Disorder Questionnaire 2007 (DCD-Q), which assesses Control During Movement (CDM), Fine Motor Skills (FMS), and General Coordination (GC) in children.

Results: Children with ADHD had more DCD symptoms overall, $F(1, 30) = 14.15, p = .001$, as well as on the CDM, $F(1, 30) = 7.72, p = .009$, FMS, $F(1, 30) = 15.57, p < .001$, and GC subscales, $F(1, 30) = 9.85, p = .004$, compared to children without ADHD. Despite no previous DCD diagnosis, 34% of children screened positive for the disorder.

Conclusions: Children with ADHD had more problems in all rated areas of motor functioning compared with children without ADHD. Many children screened positively for DCD despite not having a previous diagnosis. Findings indicate the value of using screening measures to identify children at risk of DCD. Since these deficits can cause added difficulty in the daily functioning of children with ADHD, diagnosis and treatment are especially important for this population.