

University of Windsor

Scholarship at UWindsor

UWill Discover Student Research Conference

UWill Discover 2019

Synthesis of a more biologically active anti-tumour agent

Michael A. Qaqish

University of Windsor, qaqis112@uwindsor.ca

Follow this and additional works at: <https://scholar.uwindsor.ca/uwilldiscover>

Qaqish, Michael A., "Synthesis of a more biologically active anti-tumour agent" (2024). *UWill Discover Student Research Conference*. 49.

<https://scholar.uwindsor.ca/uwilldiscover/2019/online/49>

This Event is brought to you for free and open access by the Conferences and Conference Proceedings at Scholarship at UWindsor. It has been accepted for inclusion in UWill Discover Student Research Conference by an authorized administrator of Scholarship at UWindsor. For more information, please contact scholarship@uwindsor.ca.

Synthesis of a more biologically active anti-tumour agent: Michael Qaqish, Michael Reynolds, John F. Trant

Carbohydrates fulfill many roles in biological systems including (but not limited to): acting as cellular structural support, mediating cell signaling and acting as an energy supply. They can also act as “superantigens” for the immune system when part of certain fat molecules called glycosphingolipids. These molecules are able to activate invariant Natural Killer T-Cells (iNKT cells); white blood cells that mount a dangerous, non-specific systemic immune response potentially leading to cell death. However, this same immune response (if controlled), has promise to act as a last line antiviral and/or a potential anticancer agent by potentially turning the immune system against a previously ignored virus, infection or tumour. The identification and synthesis of such immunoactive agents for clinical and mechanistic applications is of great interest in carbohydrate immunology. KRN7000 is of particular interest. It was first isolated from a marine sponge and is the most potent activator of iNKT cells. However, if almost any part of the molecule is modified, all biological activity is lost. One of the few exceptions is that the enzymatically-sensitive bond attaching the sugar to the lipid can be replaced with a stable carbon linkage. However, the synthesis is too long to be commercially useful. We wish to discuss our approach towards this molecule that attempts to streamline the approach by very subtly modifying KRN7000 in a new way. The synthesis, and its implications for biological activity, will be discussed.