

University of Windsor

## Scholarship at UWindsor

---

UWill Discover Undergraduate Conference

UWill Discover 2016

---

Mar 29th, 1:00 PM - 2:20 PM

### Community Nursing Intervention: "Victory City" arts and mental health program

Jillian Nicole Snider

*University of Windsor*, [sniderj@uwindsor.ca](mailto:sniderj@uwindsor.ca)

Follow this and additional works at: <https://scholar.uwindsor.ca/uwilldiscover>



This work is licensed under a [Creative Commons Attribution-Noncommercial 4.0 License](https://creativecommons.org/licenses/by-nc/4.0/)

---

Snider, Jillian Nicole, "Community Nursing Intervention: "Victory City" arts and mental health program" (2016). *UWill Discover Undergraduate Conference*. 11.

<https://scholar.uwindsor.ca/uwilldiscover/2016/posters2016/11>

This Event is brought to you for free and open access by the Conferences and Conference Proceedings at Scholarship at UWindsor. It has been accepted for inclusion in UWill Discover Undergraduate Conference by an authorized administrator of Scholarship at UWindsor. For more information, please contact [scholarship@uwindsor.ca](mailto:scholarship@uwindsor.ca).

# “VICTORY CITY”

## An arts and mental health community intervention for adolescents

### Description

The proposed program aims to combine the protective factors of creative expression, peer interaction, education, and access to primary health care with arts-based therapeutic interventions to improve the mental health of adolescents.

### Significance

- Mental health is severely underfunded in Canada <sup>(2)</sup>
- Suicide is a leading cause of death among adolescents <sup>(2) (4)</sup>
- 70% of mental illnesses have onset in adolescents <sup>(2) (4)</sup>
- Stigma negatively affects reception of treatment <sup>(2)</sup>



### Literature Themes

- **Best Practice for treatment:** psychiatric counselling combined with psychotropic medication <sup>(5)</sup>; only 1/3 of mentally ill adolescents complete a full clinical counseling program <sup>(1) (3) (6) (13)</sup>. In trials: MST (multisystemic therapy) proves most effective: treatment in home with whole family, focus on coping and protective factors ; MST not typically implemented to date <sup>(1)</sup> School programs more accessible, higher commitment rate: In a trial 98% of teens completed school-based programs ; 37% completed CBT; 15% completed clinic –based treatment
- **Stigma:** Parents and adolescents have negative attitudes about the use of medication in treatment <sup>(13)</sup>. Adolescents feel vulnerable in clinic settings <sup>(1) (13)</sup>
- **Education and training:** school teachers do not feel competent to screen for mental illness <sup>(3) (8) (9) (11)</sup>; art therapy can reduce grief and support trauma counseling <sup>(1) (4) (15)</sup>
- **Research Gaps:** Suicide prevention and effects on suicide rates <sup>(3) (6) (8) (9) (10) (11) (12) (14)</sup>; efficacy of treatment without symptom re-occurrence <sup>(1) (10)</sup>; lack of longitudinal studies <sup>(11) (12)</sup>; the efficacy of school-based suicide prevention programs– inconclusive results: issues with increased suicidality and social isolation <sup>(3) (8) (9)</sup>; the efficacy of art therapy in adolescent chronic mental illness <sup>(1) (4) (15)</sup>

### The Intervention – Victory City

**Upstream approach:** risk-assessment, screening, health promotion, prevention, community involvement, building capacities

- **Integrate:**
  - art therapy, mentoring and peer support into adolescent mental health programs
  - community interventions that are comprehensive and accessible
- **Build:**
  - capacities between schools, the community and the health care system
- **Evaluate:**
  - Mental health outcomes of the arts-based community program
    - Psychometric instruments: self-reported symptoms, suicidality, self-harm behaviour
    - Other indicators - hospitalizations, recurring symptoms, suicidal ideation or planning, school attendance, self-image

# BUILDING COMMUNITY CAPACITIES



## References

- (1) Asarnow, J.R., and Miranda, J. (2014). Improving Care for Depression and Suicide Risk in Adolescents: Innovative strategies for bringing treatments to community settings. *Annual Review of Clinical Psychology* Vol. 10: 275-303. DOI: 10.1146/annurev-clinpsy-032813-153742  
retrieved from: <http://www.annualreviews.org.ezproxy.uwindsor.ca/doi/full/10.1146/annurev-clinpsy-032813-153742>
- (2) Canadian Mental Health Association (CMHA). (2013). Fast Facts About Mental Illness. Retrieved from: <http://www.cmha.ca/media/fast-facts-about-mental-illness/#.VvgdCfkrLIUCiliska>, D., Dobbins, M., Hayward, S., Ploeg, J., Thomas, H., & Underwood, J., (2007) A Systematic Overview of Adolescent Suicide Prevention Programs. *Canadian Journal of Public Health / Revue Canadienne de Santé Publique*. Vol. 87, No. 5 (September/October 1996), pp. 319-324. Retrieved from: <http://www.jstor.org.ezproxy.uwindsor.ca/stable/41993812>
- (3) Ciliska, D., Dobbins, M., Hayward, S., Ploeg, J., Thomas, H., & Underwood, J., (2007) A Systematic Overview of Adolescent Suicide Prevention Programs. *Canadian Journal of Public Health / Revue Canadienne de Santé Publique*. Vol. 87, No. 5 (September/October 1996), pp. 319-324. Retrieved from: <http://www.jstor.org.ezproxy.uwindsor.ca/stable/41993812>
- (4) George, O., & Kasinathan, J. (2014). Mural art therapy for young offenders hospitalised with a mental illness. *Australasian Psychiatry*, 23(1), 49-53. Retrieved from <https://hdl.handle.net/1885/14420>.
- (5) Government of Canada. (2016). Statistics Canada: Perceived mental health by sex, by province and territory. Retrieved from: <http://www.statcan.gc.ca/tables-tableaux/sum-som/l01/cst01/health111a-eng.htm>
- (4) Karpenko, V., & Owens, J. (2013). Adolescent Psychotherapy Outcomes in Community Mental Health: How Do Symptoms Align with Target Complaints and Perceived Change?. *Community Mental Health Journal*, 49(5), 540-552. doi:10.1007/s10597-012-9515-0
- (6) Karpenko, V., & Owens, J. (2013). Adolescent Psychotherapy Outcomes in Community Mental Health: How Do Symptoms Align with Target Complaints and Perceived Change?. *Community Mental Health Journal*, 49(5), 540-552. doi:10.1007/s10597-012-9515-0

## References Continued...

- (7) Kennebeck, S., Bonin, L., (2016). Suicidal behavior in children and adolescents: Evaluation and management. Up to Date Database. Retrieved from: [http://www.uptodate.com.ezproxy.uwindsor.ca/contents/suicidal-behavior-in-children-and-adolescents-evaluation-and-management?source=search\\_result&search=adolescent+suicide+prevention&selectedTitle=1~150](http://www.uptodate.com.ezproxy.uwindsor.ca/contents/suicidal-behavior-in-children-and-adolescents-evaluation-and-management?source=search_result&search=adolescent+suicide+prevention&selectedTitle=1~150)
- (8) King, K., Strunk, C., & Sorter, M. (2011). Preliminary Effectiveness of Surviving the Teens® Suicide Prevention and Depression Awareness Program on Adolescents' Suicidality and Self-Efficacy in Performing Help-Seeking Behaviors. *Journal Of School Health*, 81(9), 581-590. doi:10.1111/j.1746-1561.2011.00630.x
- (9) Kutcher, S., Le Blanc, J., Wei, Y. (2015). Hot Idea or Hot Air: A systematic review of evidence for two widely marketed youth suicide prevention programs and recommendations for implementation. *Journal of Canadian Academic Child Adolescent Psychiatry*. 2015 Winter;24(1):5-16. Epub 2015 Mar 4. Retrieved from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4357329/>
- (10) Lichtenstein, D., Spirito, A., & Zimmermann, R. (2010). Assessing and treating co-occurring disorders in adolescents: examining typical practice of community-based mental health and substance use treatment providers. *Community Mental Health Journal*, 46(3), 252-257 6p. doi:10.1007/s10597-009-9239-y
- (11) Macleod, E., Nada-Raj, S., Beautrais, A., Shave, R., & Jordan, V. (2015). *Primary prevention of suicide and suicidal behaviour for adolescents in school settings*. Cochrane Database of Systematic Reviews 2015, Issue 12. Art. No.: CD007322. DOI: 10.1002/14651858.CD007322.pub2.
- (12) Nordentoft, M. (2007). *Prevention of suicide and attempted suicide in Denmark: Epidemiological studies of suicide and intervention studies of selected risk groups*. *Danish Medical Bulletin*, 2007 Nov;54(4):306-69. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/18208680>

## References Continued...

- (13) Oruche, U. M., Downs, S., Holloway, E., Draucker, C., & Aalsma, M. (2014). Barriers and facilitators to treatment participation by adolescents in a community mental health clinic. *Journal Of Psychiatric & Mental Health Nursing*, 21(3), 241-248. doi:10.1111/jpm.12076. Retrieved from: <http://web.a.ebscohost.com.ezproxy.uwindsor.ca/ehost/detail/detail?vid=3&sid=fffb1eb1-f3bd-4fa1-ba78-940fdff30c12%40sessionmgr4002&hid=4209&bdata=JnNpdGU9ZWwhvc3QtbGl2ZQ%3d%3d#db=a9h&AN=94727443>
- (14) Pirruccello, L. (2010). Preventing adolescent suicide: a community takes action. *Journal Of Psychosocial Nursing & Mental Health Services*, 48(5), 34-41 8p. doi:10.3928/02793695-20100303-01 Retrieved from: <http://web.b.ebscohost.com.ezproxy.uwindsor.ca/ehost/pdfviewer/pdfviewer?sid=0b65628c-4d55-4cba-b9f0-8c38932016e7%40sessionmgr120&vid=5&hid=110>
- (15) Rowe, C., Watson-Ormond, R., English, L., Rubesin, H., Marshall, A., Linton, K., . . . Eng, E. (2016). Evaluating Art Therapy to Heal the Effects of Trauma Among Refugee Youth: The Burma Art Therapy Program Evaluation. *Health Promotion Practice*. Retrieved from [http://www.academia.edu/23464311/Evaluating\\_Art\\_Therapy\\_to\\_Heal\\_the\\_Effects\\_of\\_Trauma\\_Among\\_Refugee\\_Youth\\_The\\_Burma\\_Art\\_Therapy\\_Program\\_Evaluation](http://www.academia.edu/23464311/Evaluating_Art_Therapy_to_Heal_the_Effects_of_Trauma_Among_Refugee_Youth_The_Burma_Art_Therapy_Program_Evaluation)

# THANK YOU